

BEXSERO

DOSING SCHEDULE

The doctor, pharmacist, or nurse will inject BEXSERO into the arm or leg muscle, depending on the age of the patient.

RECOMMENDED VACCINATION SCHEDULE:

| Age group at time of first dose | Primary immunization | | Booster dose (0.5 mL) |
|---|---|--------------------------------|--|
| | Primary doses <i>Each dose is 0.5 mL</i> | Interval between primary doses | |
| 2+1 dosing schedule | | | |
| Infants 2 through 5 months of age*† | 2 | ≥2 months | 1 dose required in the 2 nd year of life after an interval of ≥6 months from the last injection of the initial course.† |
| Infants 6 through 11 months of age | 2 | ≥2 months | 1 dose required in the 2 nd year of life, after an interval of ≥2 months from the last dose.† |
| Children 12 months through 23 months | 2 | ≥2 months | 1 dose required after an interval of 12 to 23 months from the second injection.† |
| Children, adolescents, and adults 2 through 25 years of age | 2 | ≥1 month | 1 dose considered in individuals at continuous risk of exposure to meningococcal disease. |

Date of initial dose: _____ / ____ / ____

* The initial course can also be given as 3 injections with ≥1 month interval between the primary doses. A booster dose should be given in the 2nd year of life after an interval of ≥6 months from the last injection of the initial course.

Vaccination against meningitis B‡ is not a part of your child’s routine vaccination schedule. Help protect your child against meningitis B‡ with BEXSERO.

Please refer to the package insert for complete dosing information.

† The need for, and timing of further doses has not yet been determined.
‡ Invasive meningococcal group B disease caused by *Neisseria meningitidis* group B strains.