

BEXSERO DOSING SCHEDULE

The doctor, pharmacist or nurse will inject BEXSERO into the arm or leg muscle depending on the age of the patient.

RECOMMENDED VACCINATION SCHEDULE:

		Primary immunization		Booster dose (0.5 mL)
Age group at time of first dose	Primary doses <i>Each dose is 0.5 mL</i>	Interval between primary doses		
NEW	Infants (Option 1) 2 through 5 months of age	2+1 dosing schedule (Option 1)		1 dose Required in the 2nd year of life after an interval of ≥ 6 months from the last injection of the initial course.
		2	≥ 2 months	
	Infants (Option 2) 2 through 5 months of age	3+1 dosing schedule (Option 2)		1 dose Required in the 2nd year of life after an interval of ≥ 6 months from the last injection of the initial course.
		3	≥ 1 month	
	Infants 6 through 11 months of age	2	≥ 2 months	1 dose Required in the 2nd year of life, after an interval of ≥ 2 months from the last dose. The need for further doses has not been established.
	Children 12 months through 23 months	2	≥ 2 months	Need not established.
	Children 2 years through 10 years of age	2	≥ 1 month	Need not established.
	Adolescents and adults 11 through 25 years of age	2	≥ 1 month	Need not established.

Date of initial dose: _____ / ____ / ____

Vaccination against MenB* is not part of your child's routine vaccination schedule. Help protect your child against MenB* with BEXSERO.